


This listing of claims will replace all prior versions, and listings, of claims in the application:

**Amendments to the Claims:**

1. (Previously presented) A method of treating urinary incontinence comprising administration of an effective amount of a compound of formula IA-IF having the following structure:



IA-IF

wherein:

the carbon atom designated \* is in the R or S configuration;

R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, each of which is optionally substituted with 1 to 3 substituents independently selected at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>;

R<sup>2</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl or C<sub>1</sub>-C<sub>6</sub> haloalkyl;

R<sup>3</sup> is H, halogen, -OR<sup>11</sup>, -S(O)R<sup>12</sup>, -S(O)<sub>n</sub>NR<sup>11</sup>R<sup>12</sup>, -CN, -C(O)R<sup>12</sup>, -C(O)NR<sup>11</sup>R<sup>12</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, -O(phenyl) or -O(benzyl), wherein each of -O(phenyl) and -O(benzyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy, or wherein R<sup>3</sup> is a C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl group, then said group is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>;

provided that for compounds of formula IA, R<sup>3</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, each of which is optionally substituted with from 1

to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>;

provided that for compounds of formula IB, R<sup>3</sup> is -O(phenyl), -O(benzyl), -OC(O)R<sup>13</sup> or -S(O)<sub>n</sub>R<sup>12</sup>, each of -O(phenyl) and -O(benzyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;

R<sup>4</sup> is H, halogen, -OR<sup>11</sup>, -S(O)<sub>n</sub>R<sup>12</sup>, -S(O)NR<sup>11</sup>R<sup>12</sup>, -CN, -C(O)R<sup>12</sup>, -C(O)NR<sup>11</sup>R<sup>12</sup>, -NR<sup>11</sup>R<sup>12</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, O(phenyl) or -O(benzyl), wherein each of -O(phenyl) and -O(benzyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl or C<sub>1</sub>-C<sub>4</sub> alkoxy and wherein R<sup>4</sup> is a C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl group, then said group is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>;

provided that for compounds of formula IC, R<sub>4</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, each of which is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>, or R<sup>5</sup> and R<sup>6</sup> or R<sup>6</sup> and R<sup>7</sup> may be -O-C(R<sup>12</sup>)<sub>2</sub>-O-;

provided that for compounds of formula ID, R<sup>4</sup> is -O(phenyl), -O(benzyl), -OC(O)R<sup>13</sup>, -NR<sup>11</sup>R<sup>12</sup> or -S(O)<sub>n</sub>R<sup>12</sup>, each of -O(phenyl) and -O(benzyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;

R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> in compounds of each of the formulae IA, IB, IC, ID, IE and IF are each independently H, halogen, -OR<sup>11</sup>, -S(O)<sub>n</sub>R<sup>12</sup>, -CN, -C(O)R<sup>12</sup>, -NR<sup>11</sup>R<sup>12</sup>, -C(O)NR<sup>11</sup>R<sup>12</sup>, -NR<sup>11</sup>C(O)R<sup>12</sup>, -NR<sup>11</sup>C(O)<sub>2</sub>R<sup>12</sup>, -NR<sup>11</sup>C(O)NR<sup>12</sup>R<sup>13</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl, wherein each of R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> is a C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or C<sub>4</sub>-C<sub>7</sub> cycloalkylalkyl group, then said group is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from C<sub>1</sub>-C<sub>3</sub> alkyl, halogen, aryl, -CN, -OR<sup>9</sup> and -NR<sup>9</sup>R<sup>10</sup>, or R<sup>5</sup> and R<sup>6</sup> or R<sup>6</sup> and R<sup>7</sup> may be -O-C(R<sup>12</sup>)<sub>2</sub>-O-;

provided that for compounds of formula IE at least one of R<sup>5</sup> or R<sup>7</sup> is fluoro, chloro, or methyl;

or  $R^7$  and  $R^6$  are each independently  $-O-C(R^{12})_2-O-$  in compounds of the formulae IE, but only where  $R^2$  is fluoro, chloro or methyl;

or  $R^7$  and  $R^6$  can independently also be  $-O-C(R^{12})_2-O-$  in compounds of the formulae IE, but only where  $R^7$  is fluoro, chloro or methyl;

$R^8$  is H, halogen, or  $OR^{11}$ , provided that for compounds of formula IF,  $R^8$  is halogen;

$R^9$  and  $R^{10}$  are each independently H,  $C_1-C_4$  alkyl,  $C_1-C_4$  haloalkyl,  $C_1-C_4$  alkoxyalkyl,  $C_3-C_6$  cycloalkyl,  $C_4-C_7$  cycloalkylalkyl,  $-C(O)R^{13}$ , phenyl or benzyl, where phenyl or benzyl is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1-C_4$  alkyl,  $C_1-C_4$  haloalkyl, or  $C_1-C_4$  alkoxy;

or  $R^9$  and  $R^{10}$  are taken together with the nitrogen to which they are attached to form piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine, or thiomorpholine;

$R^{11}$  is H,  $C_1-C_4$  alkyl,  $C_1-C_4$  haloalkyl,  $C_1-C_4$  alkoxyalkyl,  $C_3-C_6$  cycloalkyl,  $C_4-C_7$  cycloalkylalkyl,  $-C(O)R^{13}$ , phenyl or benzyl, where  $R^{11}$  is a  $C_1-C_4$  alkyl, phenyl or benzyl group, then said group is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1-C_4$  alkyl,  $C_1-C_4$  haloalkyl, or  $C_1-C_4$  alkoxy;

$R^{12}$  is H, amino,  $C_1-C_4$  alkyl,  $(C_1-C_4 \text{ alkyl})\text{amino}$ ,  $C_1-C_4$  haloalkyl,  $C_1-C_4$  alkoxyalkyl,  $C_3-C_6$  cycloalkyl,  $C_4-C_7$  cycloalkylalkyl, phenyl or benzyl, where phenyl or benzyl is optionally substituted from 1 to 3 times with a substituent selected independently from halogen, cyano,  $C_1-C_4$  alkyl,  $C_1-C_4$  haloalkyl and  $C_1-C_6$  alkoxy;

or  $R^{11}$  and  $R^{12}$  are taken together with the nitrogen to which they are attached to form piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine, or thiomorpholine; provided that only one of  $R^9$  and  $R^{10}$  or  $R^9$  and  $R^{10}$  are taken together with the nitrogen to which they are attached to form piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine, or thiomorpholine;

$R^{13}$  is  $C_1-C_4$  alkyl,  $C_1-C_4$  haloalkyl or phenyl;

$n$  is 0, 1, or 2, and;

aryl is phenyl which is optionally substituted 1-3 times with halogen, cyano,  $C_1-C_4$  alkyl,  $C_1-C_4$  haloalkyl and  $C_1-C_4$  alkoxy,

or an oxide thereof, a pharmaceutically acceptable salt thereof, a solvate thereof, or prodrug thereof.

2. (Original) A method of claim 1, wherein  $R^1$  is  $C_1-C_3$  alkyl.

3. (Original) A method of claim 2, wherein  $R^1$  is  $CH_3$ .
4. (Original) A method of claim 1, wherein  $R^2$  is H,  $C_1$ - $C_4$  alkyl or  $C_1$ - $C_6$  haloalkyl.
5. (Original) A method of claim 4, wherein  $R^2$  is H or  $CH_3$ .
6. (Original) A method of claim 1, wherein  $R^3$  is H or  $R^3$  is  $C_1$ - $C_4$  alkyl,  $C_3$ - $C_6$  cycloalkyl or  $C_4$ - $C_7$  cycloalkylalkyl, each of which is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from  $C_1$ - $C_3$  alkyl, halogen, aryl,  $-CN$ ,  $-OR^9$  and  $NR^9R^{10}$ , or  $R^3$  is  $-O(phenyl)$  or  $-O(benzyl)$  optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl, or  $C_1$ - $C_4$  alkoxy.
7. (Original) A method of claim 6, wherein  $R^3$  is methyl, ethyl, propyl, or isopropyl.
8. (Original) A method of claim 6, wherein  $R^3$  is  $-O(phenyl)$  or  $-O-CH_2-(phenyl)$ , each of which is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl, or  $C_1$ - $C_4$  alkoxy.
9. (Original) A method of claim 6, wherein  $R^3$  is H.
10. (Original) A method of claim 1, wherein  $R^4$  is H, or  $R^4$  is  $-NR^{11}R^{12}$  or  $R^4$  is  $C_1$ - $C_4$  alkyl,  $C_3$ - $C_6$  cycloalkyl or  $C_4$ - $C_7$  cycloalkylalkyl, each of which is optionally substituted, or wherein  $R^4$  is  $-O(phenyl)$  or  $-O(benzyl)$ , each of which is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl, or  $C_1$ - $C_4$  alkoxy.
11. (Original) A method of claim 10, wherein  $R^4$  is methyl, ethyl, propyl, or isopropyl.
12. (Original) A method of claim 10, wherein  $R^4$  is  $-O(phenyl)$  or  $-O(CH_2)phenyl$ , each of which is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  haloalkyl, or  $C_1$ - $C_4$  alkoxy.

13. (Original) A method of claim 10, wherein  $R^4$  is H.
14. (Original) A method of claim 1, wherein  $R^3$  and  $R^4$  are each H or wherein  $R^3$  and  $R^4$  are each halogen.
15. (Original) A method of claim 1, wherein one of  $R^3$  and  $R^4$  is H and the other is  $CH_3$ .
16. (Original) A method of claim 1, wherein  $R^5$ ,  $R^6$  and  $R^7$  are each H, halogen,  $-OR^{11}$ ,  $-NR^{11}R^{12}$ ,  $C_1$ - $C_6$  alkyl and substituted  $C_1$ - $C_6$  alkyl.
17. (Original) A method of claim 16, wherein  $R^5$ ,  $R^6$  and  $R^7$  are each H.
18. (Original) A method of claim 16, wherein one of  $R^5$  or  $R^7$  is F, Cl or Me and the other of  $R^5$  or  $R^7$  and  $R^6$  are H, halogen,  $-OR^{11}$ ,  $-NR^{11}R^{12}$ , or optionally substituted  $C_1$ - $C_6$  alkyl.
19. (Original) A method of claim 18, wherein  $R^5$  is F, Cl or Me; and  $R^7$  is H.
20. (Original) The method of claim 18, wherein  $R^5$  is F, Cl or Me; and  $R^6$  is H.
21. (Original) A method of claim 1, wherein  $R^8$  is halogen.
22. (Original) A method of claim 21, wherein  $R^8$  is fluoro.
23. (Original) A method of claim 1, wherein:  
     $R^1$  is  $C_1$ - $C_3$  alkyl;  
     $R^2$  is H,  $C_1$ - $C_4$  alkyl or  $C_1$ - $C_6$  haloalkyl;  
     $R^3$  is  $C_1$ - $C_4$  alkyl,  $C_3$ - $C_6$  cycloalkyl or  $C_4$ - $C_7$  cycloalkylalkyl, each of which is optionally substituted, or  $R^3$  is  $-O(\text{phenyl})$  or  $-O(\text{benzyl})$ , each of which is optionally substituted, or  $R^3$  is H;  $R^4$  is H,  $C_1$ - $C_4$  alkyl,  $C_3$ - $C_6$  cycloalkyl or  $C_4$ - $C_7$  cycloalkylalkyl, each of which is optionally substituted with from 1 to 3 substituents selected independently at each occurrence thereof from  $C_1$ - $C_3$  alkyl, halogen, aryl,  $-CN$ ,  $-OR^9$  and  $-NR^9R^{10}$ , or  $R^4$  is  $-NR^{11}R^{12}$ ,  $-O(\text{phenyl})$  or  $-O(\text{benzyl})$ , wherein said  $-O(\text{phenyl})$  or  $-O(\text{benzyl})$ , is optionally

substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;  
or R<sup>3</sup> and R<sup>4</sup> are each halogen;  
R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are each H, halogen, -OR<sup>11</sup>, -NR<sup>11</sup>R<sup>12</sup>, optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl, or one of R<sup>5</sup> and R<sup>7</sup> is Cl, F or Me and the other of R<sup>5</sup> and R<sup>7</sup> and R<sup>6</sup> is H, halogen, -OR<sup>11</sup>, -NR<sup>11</sup>R<sup>12</sup>, C<sub>1</sub>-C<sub>6</sub> alkyl or substituted C<sub>1</sub>-C<sub>6</sub> alkyl.

24. (Original) A method of claim 23, wherein:

R<sup>1</sup> is CH<sub>3</sub>;

R<sup>2</sup> is H or CH<sub>3</sub>;

R<sup>3</sup> is H, F, methyl, ethyl, propyl, isopropyl, -O(phenyl) or -O-CH<sub>2</sub>-(phenyl), wherein said -O(phenyl) or -O-CH<sub>2</sub>-(phenyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;

R<sup>4</sup> is H, F, methyl, ethyl, propyl, isopropyl, -O(phenyl) or -O-CH<sub>2</sub>-(phenyl), wherein said -O(phenyl) or -O-CH<sub>2</sub>-(phenyl) is optionally substituted from 1 to 3 times with a substituent selected independently at each occurrence thereof from halogen, cyano, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, or C<sub>1</sub>-C<sub>4</sub> alkoxy;

R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are each H or R<sup>5</sup> is F, Cl or Me, or one of R<sup>6</sup> or R<sup>7</sup> is H and the other of R<sup>6</sup> and R<sup>7</sup> is halogen, -OR<sup>11</sup>, -NR<sup>11</sup>R<sup>12</sup>, or optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl.

25. (Original) A method of claim 23, wherein R<sup>8</sup> is halogen.

26. (Original) A method according to claim 1, wherein the carbon atom designated \* is in the R configuration.

27. (Original) A method according to claim 1, wherein the carbon atom designated \* is in the S configuration.

28. (Original) A method comprising a mixture of stereoisomeric compounds of claim 1 wherein the carbon atom designated \* is in the S or R configuration.

29. (Previously presented) A method according to claim 1, wherein the compound is selected from the group consisting of:

2,7-dimethyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-methoxy)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
2,7-dimethyl-4-(4-fluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;  
2,7-dimethyl-4-(3-fluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-difluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
2,7-dimethyl-4-(4-fluoro-3-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro-4-fluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
2,7-dimethyl-4-(4-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;  
2,7-dimethyl-4-(3-fluoro-4-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-chloro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-chloro-3-fluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-dichloro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
7-ethyl-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-difluoro)phenyl-7-ethyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
7-fluoro-4-(4-methoxy)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
7-fluoro-4-(3-fluoro-4-methoxy)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
7-fluoro-4-(3-fluoro-4-methyl)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
7-fluoro-4-(4-chloro-3-fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-difluoro)phenyl-7-fluoro-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro)phenyl-7-fluoro-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
7-cyano-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
2-methyl-4-phenyl-7-trifluoromethyl-1,2,3,4-tetrahydroisoquinoline;  
4-phenyl-1,2,7-trimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-chloro)phenyl-1,2-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-difluoro)phenyl-1,2-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-phenyl-2,7,8-trifluoromethyl-1,2,3,4-tetrahydroisoquinoline;  
2,7-dimethyl-8-fluoro-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
2,8-dimethyl-7-fluoro-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
2,7-dimethyl-8-methoxy-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
2,7-dimethyl-8-hydroxy-4-phenyl-1,2,3,4-tetrahydroisoquinoline;

2-methyl-4-phenyl-7-trifluoromethoxy-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-difluoro)phenyl-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-fluoro-3-methyl)phenyl-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-fluoro-4-methyl)phenyl-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
7-methoxy-4-(3-methyl)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
2-methyl-7-phenoxy-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
7-(4-methoxy)phenoxy-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
7-benzyloxy-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
7-hydroxy-2-methyl-4-(3-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-fluoro-4-methyl)phenyl-7-hydroxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-fluoro-3-methyl)phenyl-7-hydroxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-difluoro)phenyl-7-hydroxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-cyano)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
2,8-dimethyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
2,8-dimethyl-4-(4-fluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-difluoro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,5-difluoro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
2,8-dimethyl-4-(3-fluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;  
2,8-dimethyl-4-(4-fluoro-3-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro-4-fluoro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-dichloro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-chloro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-chloro-3-fluoro)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
2,8-dimethyl-4-(4-methoxy)phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-cyano)phenyl-2,8-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
2,8-dimethyl-4-(4-trifluoromethyl)phenyl-1,2,3,4-tetrahydroisoquinoline;  
2,8-dimethyl-4-(4-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;  
2-methyl-8-(N-methylamino)methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
8-(hydroxy)methyl-2-methyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
2-methyl-4-phenyl-8-sulfonamide-1,2,3,4-tetrahydroisoquinoline;  
2-methyl-8-(N-methyl)sulfonamide-4-phenyl-1,2,3,4-tetrahydroisoquinoline;  
8-methoxy-2-methyl-4-(4-methyl)phenyl-1,2,3,4-tetrahydroisoquinoline;



4-(3,5-difluoro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-dichloro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-chloro-3-fluoro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro-4-fluoro)phenyl-8-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,5-difluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro-5-fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,5-difluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro-5-fluoro)phenyl-2,7-dimethyl-1,2,3,4-tetrahydroisoquinoline;  
2-methyl-4-(3,4,5-trifluoro)phenyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-fluoro-4-methyl)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-fluoro-3-methyl)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3,4-difluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-chloro-3-fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-chloro-4-fluoro)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(3-cyano)phenyl-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-acetanilide)-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
4-(4-chloro)phenyl-4-fluoro-2-methyl-1,2,3,4-tetrahydroisoquinoline;  
(3,5-difluoro)-4-phenyl-1,2,7-trimethyl-1,2,3,4-tetrahydroisoquinoline;  
(8-fluoro-2-methyl-4-phenyl-1,2,3,4-tetrahydro-7-isoquinoliny)-N-methylmethanamine;  
(2-methyl-4-phenyl-7-isoquinoliny)-N-methylmethanamine;  
N-methyl-(2-methyl-4-phenyl-7-isoquinoliny)-N-methylmethanamine;  
8-hydroxy-2-methyl-4-phenyl-1,2,3,4-tetrahydro-7-isoquinolinecarbonitrile;  
(2-methyl-4-phenyl-1,2,3,4-tetrahydro-7-isoquinoliny)methanol;  
2-ethyl-4-phenyl-1,2,3,4-tetrahydroisoquinoline; an oxide thereof, a pharmaceutically acceptable salt thereof, a solvate thereof, or prodrug thereof.

30. (Previously Presented) A method of claim 1, wherein the urinary incontinence is urge, stress, or mixed urinary incontinence.